

Vital Signs Monitoring of Wolf (*Canis lupus*) Distribution and Abundance in Denali National Park and Preserve, Central Alaska Network

2009 Report

Natural Data Series NPS/CAKN/NRDS—2009/009





Vital Signs Monitoring of Wolf (*Canis lupus*) Distribution and Abundance in Denali National Park and Preserve, Central Alaska Network

2009 Annual Report

Natural Resource Data Series NPS/CAKN/NRDS—2009/009

Thomas Meier National Park Service Denali National Park and Preserve P.O. Box 9 Denali Park, AK 99755

October 2009

U.S. Department of the Interior National Park Service Natural Resource Program Center Fort Collins, Colorado The National Park Service, Natural Resource Program Center publishes a range of reports that address natural resource topics of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report are those of the author(s) and do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the National Park Service.

This report is available from Center for Resources, Science and Learning, Denali National Park and Preserve and the Natural Resource Publications Management website (http://www.nature.nps.gov/publications/NRPM).

Please cite this publication as:

Meier, T. J., 2009. Vital signs monitoring of wolf (*Canis lupus*) distribution and abundance in Denali National Park and Preserve, Central Alaska Network. 2009 report. Natural Resource Data Series NPS/CAKN/NRDS 2009/009. National Park Service, Fort Collins, Colorado.

Contents

Page
Executive Summary
Acknowledgments
Introduction:
Measurable Objectives
Methods and Materials
Results and Discussion
Captures and Radio Telemetry
Wolf Pack Sizes and Density Estimates
Mortality5
Plans for the Coming Year6
Literature Cited
Appendix A Tables and Figures
Table 1. Early-winter density of wolves, Denali National Park and Preserve, 1986-2008.
Table 2. Late-winter density of wolves, Denali National Park and Preserve, 1986-2009 10
Table 3. Wolf pack sizes, Denali National Park and Preserve, 2004-2009
Table 4. Wolf Captures, Denali National Park and Preserve, 2003-2008
Figure 1. Wolf pack territories and population estimate for Denali National Park and Preserve, 2005
Figure 2. Wolf pack territories and population estimate for Denali National Park and Preserve, 2006
Figure 3. Wolf pack territories and population estimate for Denali National Park and Preserve, 2007
Figure 4. Wolf pack territories and population estimate for Denali National Park and Preserve, 2008

Figure 5. Wolf pack territories and population estimate for Denali National Park and Preserve, 2009	19
Figure 6. Wolf density estimates, Denali National Park and Preserve, 1986-2009	20
Figure 7. Wolf density estimates, Denali National Park and Preserve, 1986-2009, spring and fall estimates on the same plot.	21
Figure 8. Locations of VHF and GPS-collared wolves, 1986-2009	22
Figure 9. Denali National Park and Preserve, showing areas of differing wolf management.	23

Executive Summary

Wolves have been monitored with the use of radio collars in Denali National Park and Preserve since 1986. This work was conducted by the National Park Service (NPS) from 1986 to 1994, by the U. S. Geological Survey (USGS) from 1995 to 2002, and again by NPS from 2003 to 2009. A total of 137 wolves have been captured since NPS resumed wolf monitoring efforts in 2003. Between November 2008 and February 2009, 24 wolves were captured and radio collared in or near the Park and Preserve.

Each year, 15 to 20 wolf packs are monitored in or adjacent to the Park and Preserve. A total of 30 different wolves have worn GPS collars, which determine the animal's location once per day and upload the data through the ARGOS satellite system. Of 67 collared wolves that died between 2003 and 2009, 24 were killed by humans and 43 by natural causes, suggesting an increase in human-caused mortality in recent years. The estimate of wolf density in March 2009 was 3.91 wolves per 1000 square kilometers, the lowest density observed in Denali since 1987.

Acknowledgments

Nearly every aspect of wolf monitoring in Alaska is dependent on the use of aircraft. In recent years, we have relied on the skills of helicopter pilots Rick Swisher (Quicksilver Air) and Troy Cambier (Chena River Aviation), and fixed-wing pilots Dennis Miller (Caribou Air), Sandy Hamilton (Arctic Air Alaska), and Colin Milone (National Park Service). Melanie Cook of the National Park Service and Dr. Sandy Talbot of the Biological Research Division, U. S. Geological Survey, performed genetic analysis of wolf specimens. Dr. Edward DeBevec of the University of Alaska, Fairbanks, developed an online tool for decoding and managing wolf location data obtained from GPS/ARGOS telemetry collars. Telonics, Inc. (Mesa, AZ) developed radio collar designs to address unique problems of monitoring wolves in Alaska. Dr. Kimberlee Beckman of the Alaska Department of Fish and Game performed wolf necropsies and arranged for immunological testing of wolf blood specimens. John Burch and Maggie MacCluskie (NPS) reviewed this manuscript and provided additions and corrections.

Introduction

This report summarizes efforts to monitor wolves (*Canis lupus*) in Denali National Park and Preserve (DENA), Alaska, through spring 2009. Wolves occur in all three parks of the Central Alaska Monitoring Network (CAKN), Denali National Park and Preserve, Yukon-Charley Rivers National Preserve and Wrangell-St. Elias National Park and Preserve, and are one of six keystone large mammal species in interior Alaska. Wolves are of great importance to people from both consumptive and non-consumptive viewpoints, and to the ecosystem as a whole. As a top predator, wolves may play a key role in influencing ungulate populations, and as a result may influence vegetation patterns (Miller et al. 2001, Ripple and Beschta 2003). The effects of wolves on ungulate populations (Mech and Peterson 2003) may be important determinants of ungulate availability for subsistence harvest on NPS Park and Preserve lands in Alaska, and harvest by the general public on NPS Preserve lands (National Park Service 2003). Data obtained from wolf monitoring are used to assist with wolf den site protection and other aspects of the Denali Wolf-Human Conflict Management Plan (National Park Service 2007).

Wolves are a species specifically identified in the enabling legislation and management objectives of all three CAKN parks (U. S. Congress 1980). Wolves are of great importance to park visitors because of the unique opportunities to view wolves in Alaskan parks. While the primary objectives of monitoring is to track the distribution and abundance of wolves, a variety of accessory data is obtained in the monitoring process. This information is likely to have great value for wildlife management and research. The body of data on wolf populations in Alaska parks is of great value in developing scientific models of predator/prey systems. In heavily visited portions of the parks, managers may want to know the locations of active wolf dens and rendezvous sites so that they can be protected from disturbance. When intensive wolf harvest or wolf control take place near parks, it is vital to know the patterns of travel of park packs, in order to determine whether they are being significantly impacted by activities outside the parks. Data on the genetic and morphological characteristics of wolves, obtained in the course of wolf capture, will be important in evaluating long-term changes in wolf populations in Alaska.

Parkwide monitoring of wolves in Denali National Park and Preserve was initiated by Resource Management Ranger John Dalle-Molle in 1986, with principal investigators L. David Mech and Layne Adams. Field work between 1986 and 1994 was performed by John Burch and Tom Meier. From 1995 through 2002, Layne Adams, now with USGS, conducted wolf monitoring efforts. Since 2003, John Burch and Tom Meier have again conducted the field work.

Measurable Objectives

- Locate non-radiocollared wolf packs using Park and Preserve lands by snow tracking.
- Capture and radio-collar 1-3 individuals in each wolf pack identified in the study area.
- Determine the demography (numbers, colors, age structure) of monitored wolf packs.
- Obtain morphological measurements from captured wolves.
- Obtain genotypic data (mitochondrial and microsatellite DNA) from captured wolves.
- Obtain immunological (disease exposure) data from captured wolves.
- Determine pack size for each collared pack in fall (early winter) and spring (late winter).

- Define the mosaic of wolf home ranges (population area) for estimating wolf densities.
- Perform annual capture efforts to maintain coverage of radio collars in the population.
- Detect pack extinction and pack formation events in the population.
- Detect changes in wolf density over time.
- Detect changes in wolf pack sizes over time.
- Detect changes in wolf home ranges over time.
- Detect changes in the morphological, immunological, and genetic makeup of the wolf population over time.

Methods and Materials

Methods of wolf monitoring used in DENA followed the Wolf Monitoring Protocol for Denali National Park and Preserve, Yukon-Charley Rivers National Preserve and Wrangell-St. Elias National Park and Preserve, Alaska (Meier et al. 2009). An exception to this is the determination of wolf pack territories (Figure 1-5, see Appendix). Wolf pack territories were not rigorously calculated using the 95% of locations that would produce the smallest home range. Appropriate software must be found or developed in order to automate this task. The present wolf pack territories were produced by hand removing selected points that were thought by the author to represent extraterritorial forays or pre-dispersal movements by the collared animals.

Results and Discussion

Captures and Radio Telemetry

Twenty-four wolves from 17 packs were captured and radio-collared in winter 2008-2009 (Table 4 in the Appendix). Four new pairs of wolves were discovered by snow tracking in winter 2008-2009. The Iron Creek pair, southwest of Wonder Lake, were seen but not captured. Collars were placed on three new pairs; Moose Creek (in the area formerly occupied by the Turtle Hill Pack), Nenana River (in the area formerly occupied by the Mount Margaret Pack, which had moved farther north), and Otter Lake (in the area formerly occupied by the Castle Rocks III Pack) (Table 4). Collared wolves from the Somber, Bearpaw, Totek Hills, Chitsia, and Toklat Springs Packs dispersed from their territories and may provide radio collars in newly-formed groups in winter 2009-2010.

Morphological data, including sex, age and color, and blood and tissue samples, for genetics and disease analysis, were gathered from captured wolves. Morphologic data is presented in Table 4 (see appendix). Genetics results are being analyzed by biologists at United States Geological Survey (USGS) and University of California Los Angeles (UCLA). Results from immunological samples show that wolves living in or near Denali have been infested with the dog louse *Trichodectes canis* and also with another coat abnormality of unknown origin (Beckmen et al. 2009, Wolstad et al. 2009). The Alaska Department of Fish and Game and Washington Animal Disease Laboratory are testing Denali wolf serum for brucella, toxoplasma, neospora, tularemia, canine adenovirus, coronavirus, canine parvovirus (CPV), canine parainfluenza, and canine herpes virus. Immunological surveys of wolves in interior Alaska have revealed exposure to a number of these diseases but have not detected evidence of serious population effects of disease (Mech et al. 1998, Zarnke and Ballard 1987). One disease that has the potential to seriously

affect wolf pup survival is CPV. Immunological studies of Denali wolves have revealed a rate of exposure to CPV as high as 50% in some years, among wolves that were captured and blood sampled (R. Zarnke, pers. comm.).

From 1 May 2008 through 1 May 2009 collared wolves were located approximately twice per month by aircraft. Fifty different radio-collared wolves from 21 packs were monitored, resulting in 611 locations of collared wolves. In addition, 1,592 locations were obtained from 13 wolves that wore GPS/ARGOS collars for some or all of this period. The Telonics GPS collars used on these wolves obtain one location each day and store the location coordinates within the collar. The data is uploaded weekly through the ARGOS satellite system, and also remains stored within the collar so that all data can be uploaded when the collar is retrieved. Since 2003, more than 8,600 wolf locations have been obtained from GPS/ARGOS collars. The locations of all collared wolves in this study's 23-year history are plotted on Figure 8 in the Appendix.

Wolf Pack Sizes and Density Estimates

Aircraft surveys in spring 2009 observed sixty-five wolves, 29 of them radio-collared. These wolves were found in 16 packs covering an area of 16,607 square kilometers, mostly within the boundaries of Denali National Park and Preserve north of the Alaska Range (Figure 5 and Table 3, in Appendix). This produced a density estimate of 3.91 wolves per 1000 square kilometers, the lowest estimated wolf density in the park/preserve since 1987 (Table 1, 2, Figures 6 and 7 in Appendix). No obvious explanation for this low density is apparent, and it is expected that wolf numbers will rebound in winter 2009-2010.

The present method of determining wolf density involves the use of minimum convex polygons to estimate individual wolf pack territories, and combining a number of territories into a larger, not necessarily convex polygon representing the population. In implementing this method, subjective decisions are by made NPS wildlife biologists to exclude forays by wolf packs outside of their usual range, so that the population area is not inflated by the inclusion of areas that are actually occupied by other, uncollared wolf packs. CAKN personnel are developing methods using kernel estimators (White and Garrott 1990) that might provide a more objective estimate of pack territory sizes and wolf densities. Dispersing or lone wolves were not included in population size or density estimates.

Mortality

Seventeen radio-collared wolves died between 1 May 2008 and 1 May 2009. Five were legally trapped or snared outside of the Park/Preserve. Eleven died of natural causes. One wolf died of capture-related causes when its nose and mouth melted through thin ice at the release site, causing it to drown.

Table 4 (see Appendix) summarizes the fates of wolves captured and radio-collared between March 2003 and March 2009. Of 68 radio-collared wolves that were captured during this period and subsequently died, 24 (35%) were killed by humans. This suggests an increase in human-caused mortality in the DENA wolf population. During the period 1986-1994, only 8 of 58 (14%) of mortalities of radio-collared wolves were human-caused (Mech et al. 1998). Although

a buffer zone created in 2001 prevents legal harvest of wolves in some areas adjacent to the park (Figure 9 in Appendix), harvest pressure appears to have increased in easily accessible areas where trapping and hunting of wolves is legal. For both early and recent periods, subsistence harvest of wolves within Park/Preserve boundaries by qualified rural residents (National Park Service 2003) represents 10% of human harvest of radio-collared wolves (1.4% and 3.5% of total collared wolf mortality, respectively).

Intraspecific strife (the killing of wolves by members of neighboring wolf packs) probably remains the leading cause of wolf mortality in DENA (Mech et al. 1998), but many carcasses are consumed or decomposed before they can be investigated, so that only 30% of natural mortality from this sample was documented as wolf-caused, while most mortalities were classified as of unknown natural cause. It is likely that many of these mortalities were also wolf-caused.

Plans for the Coming Year

In 2009-2010, we plan to maintain contact with approximately 15-18 wolf packs inside or partly inside Denali National Park and Preserve. Collars will be maintained on 2 members of each pack if possible. Six additional Gen 4 Telonics GPS/ARGOS collars have been purchased and three Gen 3 GPS/ARGOS collars refurbished. Nine Telonics VHF collars have been refurbished to a new configuration (Mod-515) with multiple batteries designed for greater durability. Monitoring efforts will continue, with wolves being located about twice per month, with extra monitoring flights in spring and fall to document pack sizes and pup production.

Literature Cited

- Beckmen, K.B., G.D. Bossart, and K. Burek. 20009. A novel dermatopathy in Alaskan gray wolves (*Canis lupus*). Poster presentation at the Northwest Sectional Meeting of the Wildlife Society, Fairbanks, AK.
- Mech, L. D., L. G. Adams. T. J. Meier, J. W. Burch, and B. W. Dale. 1998. <u>The Wolves of Denali</u>. University of Minnesota Press, MN. 227 pages.
- Mech, L. D., and R. O. Peterson. 2003. Wolf-prey relations. Pp. 131-160 *In* Mech, L.D., and L. Boitani, eds., Wolves: Behavior, Ecology, and Conservation. University of Chicago Press, IL. 448 pp.
- Meier, T. J., Burch, J. W., Wilder, D., Cook, M. 2009. Wolf monitoring protocol for Denali National Park and Preserve, Yukon-Charley Rivers National Preserve and Wrangell-St. Elias National Park and Preserve, Alaska. Natural Resource Report NPS/CAMN/NRR—20XX/XXX. National Park Service, Fort Collins, CO.
- Miller, B., B. Dugelby, D. Foreman, C. Martinez del Rio, R. Noss, M. Phillips, R. Reading, M. E. Soule', J. Terborgh, and L. Willcox. 2001. The importance of large carnivores to healthy ecosystems. Endangered Species Update 18: 202-210.
- National Park Service, 2003. Subsistence management plan, Denali National Park and Preserve.
- National Park Service, 2007. Wolf-human conflict management plan, Denali National Park and Preserve. Denali National Park and Preserve, AK. 85 pp.
- Ripple, W.J., and R. L. Beschta. 2003. Wolf reintroduction, predation risk, and cottonwood recovery in Yellowstone National park. Forest Ecology and Management 184: 299-313.
- U. S. Congress. 1980. Alaska National Interest Lands Conservation Act, PUBLIC LAW 96-487-DEC. 2, 1980. 94 STAT. 2371. http://www.r7.fws.gov/asm/anilca/toc.html
- White, G. C., and R. A. Garrott. 1990. Analysis of wildlife radio-tracking data. Academic Press, San Diego, CA.
- Wolstad, T., K. Beckmen, C. Gardner, F. Huettmann, and K. Hundertmark. 2009. Lousy wolves: Distribution and habitat assessment of Trichodectes canis, an invasive ectoparasite of Alaskan gray wolves.
- Zarnke, R.L., and W.B. Ballard. 1987. Serological survey for selected microbial pathogens of wolves in Alaska, 1975-1982. J. Wildl. Dis. 23:77-85.
- Zarnke, R.L., J. Evermann, J.M.VerHoef, M.E. McNay, R.D. Boertje, C.L. Gardner, L.G. Adams, B.W. Dale, and J.W. Burch. 2001. Serologic survey for canine coronavirus in wolves from Alaska. J. Wildl. Dis. 37:740-745.

Appendix A Tables and Figures

Tables

- Table 1. Early-winter density of wolves, Denali National Park and Preserve, 1986-2008.
- Table 2. Late-winter density of wolves, Denali National Park and Preserve, 1986-2009.
- Table 3. Wolf pack sizes, Denali National Park and Preserve, 2004-2009.
- Table 4. Wolf captures, Denali National Park and Preserve, 2003-2008.

Figures

- Figure 1. Wolf pack territories and population estimate for Denali National Park and Preserve, 2005.
- Figure 2. Wolf pack territories and population estimate for Denali National Park and Preserve, 2006.
- Figure 3. Wolf pack territories and population estimate for Denali National Park and Preserve, 2007.
- Figure 4. Wolf pack territories and population estimate for Denali National Park and Preserve, 2008.
- Figure 5. Wolf pack territories and population estimate for Denali National Park and Preserve, 2009.
- Figure 6. Wolf density estimates, Denali National Park and Preserve, 1986-2009.
- Figure 7. Wolf density estimates, Denali National Park and Preserve, 1986-2009, spring and fall estimates on the same plot.
- Figure 8. Locations of VHF- and GPS-collared wolves, 1986-2009.
- Figure 9. Map of Denali National Park and Preserve showing zones of wolf management.

Table 1. Early-winter (fall) density of wolves, Denali National Park and Preserve, 1986-2008.

YEAR	NUMBER OF PACKS MONITORED	TOTAL WOLVES IN PACKS MONITORED	COMBINED AREA OF MONITORED PACKS (KM ²)	ESTIMATED DENSITY: WOLVES / 1000 KM ²	POPULATION ESTIMATE INSIDE THE PARK
1986	4	22	8,180	2.7	46
1987	9	70	13,150	5.3	92
1988	14	121	14,670	8.2	142
1989	11	127	15,240	8.3	144
1990	11	136	13,930	9.8	169
1991	13	137	14,275	9.6	166
1992	15	120	13,620	8.8	152
1993	12	93	9,900	9.4	162
1994	12	72	11,145	6.5	112
1995	11	80	12,045	6.6	115
1996	11	104	12,776	8.1	141
1997	12	75	12,808	5.9	101
1998	12	68	12,578	5.4	93
1999	15	80	12,699	6.3	109
2000	18	112	14,554	7.7	133
2001	18	91	13,802	6.6	114
2002	14	86	12,226	7.0	121
2003	11	84	11,682	7.2	124
2004	14	78	14,630	5.3	92
2005	15	106	15,367	6.9	119
2006	17	111	17,439	6.4	110
2007	20	147	17,757	8.3	143
2008	14	86	16,607	5.2	89

Table 2. Late-winter (spring) density of wolves, Denali National Park and Preserve, 1986-2009.

		TOTAL	COMBINED	ESTIMATED	POPULATION
	NUMBER	WOLVES	AREA OF	DENSITY:	ESTIMATE
	OF PACKS	IN PACKS	MONITORED	WOLVES /	INSIDE THE
YEAR	MONITORED	MONITORED	PACKS (KM²)	1000 KM ²	PARK
1986	4	26	7,380	3.5	61
1987	8	37	12,125	3.1	53
1988	14	69	15,355	4.5	78
1989	13	98	16,810	5.8	101
1990	10	106	13,930	7.6	131
1991	13	111	14,275	7.8	134
1992	15	103	13,620	7.6	131
1993	12	68	9,900	6.9	119
1994	10	61	11,145	5.5	95
1995	12	59	12,120	4.9	84
1996	11	69	12,640	5.5	94
1997	11	78	13,080	6.0	103
1998	12	61	13,121	4.6	80
1999	13	69	12,699	5.4	94
2000	17	71	14,378	4.9	85
2001	16	87	13,802	6.3	109
2002	15	73	13,026	5.6	97
2003	18	75	11,682	6.4	111
2004	14	78	16,061	4.9	84
2005	15	66	14,630	4.5	78
2006	15	103	15,367	6.7	116
2007	16	93	17,439	5.3	92
2008	20	99	17,757	5.6	96
2009	16	65	16,607	3.9	68

Table 3. Wolf pack sizes, Denali National Park and Preserve, 2004-2009.

PACK	2004		2005		2006		2007		2008		2009
	spring	fall	spring								
100 MILE	8	0									
BEARPAW	2	5	3	6	6	10	8	10	9	5	5
BOOT LAKE								1	(2)*	2	4
CASTLE ROCKS	1	1	1	2	2	6	5	7	7	2	0
CHITSIA	4	4	4	8	6	11	9	11	5	7	1
EAST FORK	7	12	6	14	14	15	15	15	11	16	11
GRANT CREEK	2	5	4	6	6	13	5	5	3	6	6
HAUKE						4	4	4	3	0	0
HERRON	8	8	8	3	3	0					
HOT SLOUGH							2	7	7	8	6
HULT			1								
IRON CREEK											2
KANTISHNA RIVER	5	2	2	7	7	1	1	1	2	5	2
MCKINLEY RIVER	2	6	4	2	2	6	5	10	2	0	0
MCKINLEY SLOUGH	7	8	8	10	10	8	8	15	15	14	11
MCLEOD 2							4	6	2	0	
MOOSE CREEK											2
MT MARGARET	9	7	6	10	9	11	6	7	3	2	2
MUDDY	1										
NENANA RIVER											2
OTTER LAKE											2
PINTO				6	6	5	3	10	4	0	0
SAVAGE								2	2	6	0
SOMBER						7	6	11	8	8	4
STARR LAKE	7	7	7	9	9	3	3	6	4	3	3
STRAIGHTAWAY	10	4	4	8	8	0					
TOKLAT SPRINGS	5	7	6	12	12	8	6	9	6	(5)*	(1)*
TONZONA								(1)*	(2)*	2	2
TOTEK HILLS								6	6	(4)*	(3)*
TURTLE HILL		2	2	3	3	3	3	4	0		
TOTAL WOLVES	78	78	66	106	103	111	93	147	99	86	65

^{*}Peripheral packs, with sizes in parentheses, were not included in the density estimate that year.

Table 4. Wolf captures, Denali National Park and Preserve, 2003-2008.

WOLF	DATE	RECAP	SEX	AGE	COLOR	PACK	STATUS / FATE
0087	03/16/03	Yes	F	Adult ~8	Dark Gray	Kantishna River	Recaptured, recollared
0316	03/16/03	No	М	Adult ~4	Gray	Castle Rocks 2	Killed by wolves
9755	03/16/03	Yes	F	Adult ~6	Light Gray	Pinto	Shot west of Healy
9974	03/16/03	Yes	М	Adult ~2	Dark Gray	Death Valley	Died, unknown natural causes
0092	03/17/03	Yes	F	Adult ~5	Gray	Mount Margaret	Killed by wolves
0317	03/17/03	No	F	Adult ~3	Black	Castle Rocks 2	Snared near Lake Minchumina
0079	03/18/03	Yes	М	Adult ~6	Blue	Muddy River	Died, unknown natural causes
0089	03/18/03	Yes	М	Adult ~6	Gray	McKinley Slough	Missing
0104	10/26/03	Yes	М	Adult ~4	Gray	Mount Margaret	Snared west of Healy
0318	10/26/03	No	F	Adult ~4	Gray	East Fork	Snared west of Healy
0103	10/27/03	Yes	F	Adult ~4	Gray	Grant Creek	Collar failed
0319	10/27/03	No	М	Adult ~2	Gray	Death Valley	Trapped west of Healy
0320	10/28/03	No	F	Yearling	Gray	McKinley Slough	Missing
0321	10/28/03	No	F	Adult ~4	Gray	Straightaway	Starved
0322	10/28/03	No	F	Yearling	Black	Kantishna River	Missing
0401	02/18/04	No	F	Yearling	Gray	Muddy River	Shot west of Healy
0402	02/18/04	No	F	Adult ~5	Blue	Chitsia	Recaptured, recollared
0403	03/06/04	No	F	Adult ~4	Gray	Turtle Hill	Collar failed
0102	03/07/04	Yes	F	Adult ~6	Light Gray	100 Mile	Died, probably starved
0404	03/07/04	No	М	Adult ~5	Light Gray	100 Mile	Starved
0405	03/07/04	No	F	Yearling	Gray	100 Mile	Died of unknown causes
0406	03/07/04	No	F	Adult ~5	Gray	Herron	Collar chewed off
0407	03/07/04	No	М	Adult ~2	Gray	Herron	Collar chewed off
0408	03/08/04	No	F	Yearling	Gray	Mount Margaret	Recaptured, recollared
0409	03/08/04	No	F	Adult ~4	White	Toklat Springs	Recaptured, recollared
0410	03/08/04	No	М	Yearling	Gray	Toklat Springs	Shot on lower Toklat River
0411	03/08/04	No	F	Yearling	Gray	Chitsia	Recaptured, recollared
0412	03/08/04	No	F	Adult ~4	Gray	McKinley River	Died, unknown natural causes
0413	03/08/04	No	М	Pup	Gray	Castle Rocks 2	Died, unknown natural causes
0087	03/09/04	Yes	F	Adult ~9	Blue	Kantishna River	Starved
0414	03/09/04	No	М	Adult ~4	Black	Bearpaw	Recaptured, recollared
0415	03/09/04	No	F	Adult ~3	Gray	Bearpaw	Recaptured, recollared
0416	03/09/04	No	F	Pup	Gray	Starr Lake	Missing
0411	03/04/05	Yes	F	Adult ~2	Gray	Chitsia	Collar failed
0414	03/04/05	Yes	М	Adult ~5	Black	Bearpaw	Recaptured, recollared
0501	03/04/05	No	М	Adult ~3	Gray	Mount Margaret	Recaptured, recollared
0502	03/04/05	No	F	Yearling	Gray	Mount Margaret	Died, unknown natural causes
0503	03/04/05	No	М	Adult ~2	Gray	Chitsia	Recaptured, recollared
0504	03/04/05	No	М	Adult ~8	White	Toklat Springs	Shot on lower Toklat River
0505	03/05/05	No	М	Adult ~3	Gray	McKinley River	Recaptured, recollared
0101	03/06/05	Yes	F	Adult ~6	Gray	Kantishna River	Killed by wolves
0107	03/06/05	Yes	F	Adult ~6	Gray	McKinley Slough	Killed by wolves
0506	03/06/05	No	М	Adult ~3	Black	Lone (Hult Creek)	Starved
0507	03/06/05	No	М	Adult ~3	Black	Kantishna River	Killed by wolves
0508	03/07/05	No	F	Adult ~4	Gray	Turtle Hill	Killed by wolves

Table 4. (Continued). Wolf captures, Denali National Park and Preserve, 2003-2008.

WOLF	DATE	RECAP	SEX	AGE	COLOR	al Park and Preserv	STATUS / FATE
0408		Yes	F				
	02/01/06	No	M	Adult ~3	Gray	Mount Margaret	Died from blunt trauma
0601	02/01/06			Adult ~2	Black	Turtle Hill	Trapped west of Healy
0403	02/08/06	Yes	F	Adult ~6	Light Gray	Turtle Hill	Recaptured, recollared
0602	02/09/06	No	F	Yearling	Gray	Toklat Springs	Shot on lower Teklanika River
0605	02/23/06	No	F	Pup	Black	East Fork	Shot near Cantwell
0606	02/23/06	No	M	Adult ~3	Gray	Pinto	Died of unknown causes
0607	02/23/06	No	F	Adult ~5	Gray	Pinto	Shot west of Healy
0103	02/24/06	Yes	F	Adult ~6	Gray	Grant Creek	Killed by wolves
0215	02/24/06	Yes	M	Adult ~6	Gray	Grant Creek	Killed by wolves
0214	02/28/06	Yes	F	Adult ~7	Blue	McKinley River	Killed by wolves
0505	02/28/06	Yes	M	Adult ~4	Gray	McKinley River	Recaptured, recollared
0608	02/28/06	No	F	Adult ~4	Silver	Starr Lake	Drowned
0609	02/28/06	No	F	Adult ~3	Black	Starr Lake	Recaptured, recollared
0610	03/01/06	No	F	Adult ~4	Gray	Somber	Recaptured, recollared
0611	03/01/06	No	F	Yearling	Gray	Somber-Tonzona	Still being monitored
0612	03/01/06	No	F	Adult ~4	Gray	Castle Rocks 3	Recaptured, recollared
0613	03/01/06	No	M	Adult ~3	Blond	Castle Rocks 3	Missing
0614	03/01/06	No	M	Yearling	Gray	Toklat Springs	Trapped south of Healy
0615	03/11/06	No	M	Adult ~7	White	Straightaway	Killed by wolves
0616	04/17/06	No	M	Pup	Gray	Kantishna River	Trapped in Minto area
0617	04/17/06	No	M	Pup	Black	Kantishna River	Recaptured, recollared
0618	04/17/06	No	F	Adult ~2	Gray	East Fork	Recaptured, recollared
0503	12/11/06	Yes	М	Adult ~4	Gray	Chitsia	Still being monitored
0620	12/11/06	No	F	Yearling	Gray	Chitsia	Trapped north of park boundary
0621	12/11/06	No	F	Yearling	Gray	Toklat Springs	Trapped west of Healy
0622	12/11/06	No	М	Yearling	Gray	Grant Creek	Snared east of park boundary
0701	02/14/07	No	М	Adult ~2	Gray	McKinley Slough	Missing
0702	02/14/07	No	М	Adult ~3	Gray	McKinley Slough	Recaptured, recollared
0703	02/14/07	No	F	Adult ~8	White	Hauke	Killed by wolves
0704	02/14/07	No	М	Adult ~4	Gray	Hauke	Recaptured, recollared
0402	02/27/07	Yes	F	Adult ~8	Gray	Chitsia	Died, unknown natural causes
0409	02/27/07	Yes	F	Adult ~6	White	Pinto	Trapped west of Healy
0705	03/01/07	No	F	Yearling	Light Gray	McLeod 2	Killed by wolves
0706	03/01/07	No	F	Adult ~7	Gray	McLeod 2	Died, unknown natural causes
0707	03/01/07	No	F	Adult ~2	Gray	Lone (Boot Lake)	Still being monitored
0505	03/03/07	Yes	М	Adult ~5	Gray	McKinley River	Died ,unknown natural causes
0708	03/03/07	No	F	Pup	Gray	Somber	Still being monitored
0709	03/03/07	No	F	Pup	Gray	Bearpaw	Collar found near Manley
0710	03/04/07	No	F	Pup	Gray	Somber	Shot west of park boundary
0711	03/04/07	No	M	Pup	Gray	Totek Hills	Still being monitored
0712	03/28/07	No	F	Yearling	Black	East Fork/Savage	Trapped south of Healy
0713	03/28/07	No	M	Pup	Gray	Totek Hills	Died, unknown natural causes
0610	03/29/07	Yes	F	Adult ~6	Gray	Somber	Died, unknown natural causes
0612	03/29/07	Yes	F	Adult ~5	Blue	Castle Rocks 3	Died, unknown natural causes
0714	03/29/07	No	F	Adult ~2	Gray	Hot Slough	Killed by wolves
0715	03/29/07	No	M	Adult ~2	Gray	Hot Slough	Died, unknown natural causes
0,10	00,20/01	110	171	/ wuit ~Z	Jiuy	. iot olougii	2.34, diminowii naturai causes

Table 4. (Continued). Wolf captures, Denali National Park and Preserve, 2003-2008.

	Table 4. (Continued). Wolf captures, Denali National Park and Preserve, 2003-2008.									
WOLF	DATE	RECAP	SEX	AGE	COLOR	PACK	STATUS / FATE			
0403	11/27/07	Yes	F	Adult ~8	White	Turtle Hill	Died, unknown natural causes			
0702	11/27/07	Yes	М	Adult ~4	Gray	McKinley Slough	Still being monitored			
0717	11/27/07	No	М	Adult ~5	Silver	East Fork	Still being monitored			
0718	11/27/07	No	F	Adult ~5	Light Gray	McKinley Slough	Died, unknown natural causes			
0719	11/27/07	No	F	Adult ~2	Gray	Grant Creek	Still being monitored			
0720	11/28/07	No	М	Adult ~3	Gray	McLeod 2	Trapped north of park boundary			
0721	11/28/07	No	М	Yearling	Gray	Hauke	Dispersed north			
0722	11/28/07	No	F	Adult ~3	Black	Kantishna River	Recaptured, recollared			
0723	11/29/07	No	М	Adult ~3	Gray	Toklat Springs	Still being monitored			
0724	11/29/07	No	F	Pup	Gray	Totek Hills	Trapped north of park boundary			
0414	03/02/08	Yes	М	Adult ~8	Silver	Bearpaw	Died, unknown natural causes			
0415	03/02/08	Yes	F	Adult ~7	Gray	Bearpaw	Still being monitored			
0618	03/02/08	Yes	F	Adult ~4	Gray	East Fork	Still being monitored			
0801	03/02/08	No	М	Adult ~2	Gray	Kabena	Pack moved north out of area			
0704	03/03/08	Yes	М	Adult ~5	Gray	Hauke	Killed by wolves after capture			
0802	03/03/08	No	М	Yearling	Gray	Castle Rocks 3	Died, unknown natural causes			
0803	03/03/08	No	F	Yearling	Gray	Unk/Fish Camp	Died, unknown natural causes			
0609	03/04/08	Yes	F	Adult ~5	Black	Starr Lake	Still being monitored			
0804	03/04/08	No	М	Adult ~2	Gray	Starr Lake	Still being monitored			
0805	03/04/08	No	F	Pup	Gray	Hot Slough	Still being monitored			
0806	03/04/08	No	М	Adult ~2	Gray	Castle Rocks 3	Dispersed, shot in wolf control			
0807	04/02/08	No	М	Adult ~3	Gray	Savage	Not collared, last seen 12/08			
0501	11/01/08	Yes	М	Adult ~2	Gray	Mount Margaret	Trapped west of Healy			
0810	11/01/08	No	F	Adult ~2	Gray	Toklat Springs	Trapped north of park boundary			
0811	11/01/08	No	М	Adult ~3	Gray	Grant Creek	Still being monitored			
0812	11/01/08	No	F	Adult ~5	Light Gray	McKinley Slough	Still being monitored			
0813	11/01/08	No	F	Adult ~7	Black	Mount Margaret	Still being monitored			
0814	11/02/08	No	М	Adult ~7	Dark Gray	Totek Hills	Still being monitored			
0815	11/02/08	No	F	Yearling	Yellow gray	Hot Slough	Trapped south of Minchumina			
0816	11/02/08	No	F	Adult ~3	Gray	McKinley Slough	Melted ice, drowned at capture			
0817	11/03/08	No	М	Yearling	Dark Gray	Somber	Died, unknown natural causes			
0818	11/03/08	No	М	Adult ~4	Gray	Tonzona	Still being monitored			
0819	11/04/08	No	М	Adult ~4	Gray	Chitsia	Died, unknown natural causes			
0617	02/21/09	Yes	М	Adult ~3	Black	Kantishna River	Still being monitored			
0722	02/21/09	Yes	F	Adult ~4	Black	Kantishna River	Still being monitored			
0901	02/21/09	No	F	Yearling	Gray	Hot Slough	Still being monitored			
0902	02/22/09	No	М	Adult ~2	Black	Bearpaw	Still being monitored			
0903	02/22/09	No	F	Yearling	Gray	Somber	Still being monitored			
0904	02/22/09	No	М	Adult ~2	Gray	Boot Lake	Still being monitored			
0905	02/23/09	No	F	Yearling	Gray	Nenana River	Still being monitored			
0906	02/23/09	No	М	Adult ~3	Black	Mount Margaret	Still being monitored			
0907	02/23/09	No	F	Adult ~2	Blue	Otter Lake	Still being monitored			
0908	02/23/09	No	F	Adult ~2	Black	Otter Lake	Still being monitored			
0909	02/23/09	No	F	Adult ~2	Gray	Moose Creek	Still being monitored			
0910	02/23/09	No	М	Adult ~2	Gray	Moose Creek	Died, unknown natural causes			
0911	02/24/09	No	М	Adult ~3	Gray	Nenana River	Still being monitored			




Figure 1. Wolf pack territories and population estimate for Denali National Park and Preserve, 2005.




Figure 2. Wolf pack territories and population estimate for Denali National Park and Preserve, 2006.




Figure 3. Wolf pack territories and population estimate for Denali National Park and Preserve, 2007




Figure 4. Wolf pack territories and population estimate for Denali National Park and Preserve, 2008.




Figure 5. Wolf pack territories and population estimate for Denali National Park and Preserve, 2009.




Figure 6. Wolf density estimates, Denali National Park and Preserve, 1986-2009.




Figure 7. Wolf density estimates, Denali National Park and Preserve, 1986-2009, spring and fall estimates on the same plot.




Figure 8. Locations of VHF- and GPS-collared wolves, 1986-2009.




Figure 9. Denali National Park and Preserve, showing areas of differing wolf management.